



PRE AND POST CARE INSTRUCTIONS BHRT PELLETT TREATMENT MALE

Pre Treatment

- Avoid blood thinning agents such as NSAIDS (non-steroidal anti-inflammatory drugs like Aspirin, Motrin, Advil, Ibuprofen or any other non-Tylenol, non-acetaminophen product) or Fish Oil capsules for 1 week prior to treatment.

Post Treatment

- Your insertion site has been covered with 2 layers of bandages. Remove the outer pressure bandage the morning after your insertion.
- You may remove the inner layer of steri-strips after 5 days or let them fall off naturally.
- Do not take baths, hot tub or go in a swimming pool for 7 days. You may shower but do not scrub the site for 7 days.
- If your insertion was above the belt line, no major back muscle exercises for the next 7 days that would stretch/stress/twist the insertion site including tennis, golf and side bends.
- If your insertion was below the belt line, no major gluteal exercises for 7 days including running and squats.
- You may have slight swelling for 1-3 days.
- Bruising, swelling or redness of the insertion site may last from a few days to 2-3 weeks.
- There may be pinkish or bloody discharge.
- If you start bleeding, apply firm pressure for 5 minutes.

**We want you to be completely happy with your visit to Eterna!
Call the office with any questions or concerns 253.268.3400 or after hours at
253.249.6979.**

**If you are pleased with the service you received today, we invite you to post a review on
Yelp or Google.**